



## Simulating the Ultrafast Excited State Dynamics of Cyclopentadiene and 1,2,3,4-tetramethylcyclopentadiene

Kuhlman, Thomas Scheby; Glover, Will J.; Mori, Toshifumi; Møller, Klaus Braagaard; Martinez, Todd J.

*Publication date:*  
2012

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Kuhlman, T. S., Glover, W. J., Mori, T., Møller, K. B., & Martinez, T. J. (2012). *Simulating the Ultrafast Excited State Dynamics of Cyclopentadiene and 1,2,3,4-tetramethylcyclopentadiene*. Poster session presented at CECAM conference, Birmingham, United Kingdom.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Simulating the Ultrafast Excited State Dynamics of Cyclopentadiene and 1,2,3,4-tetramethylcyclopentadiene

Thomas S. Kuhlman<sup>†</sup>, Will J. Glover<sup>‡</sup>, Toshifumi Mori<sup>‡</sup>, Klaus B. Møller<sup>†</sup>, and Todd J. Martínez<sup>‡</sup>

<sup>†</sup> Department of Chemistry, Technical University of Denmark, DTU Kemitorvet 207, DK-2800 Kgs. Lyngby, Denmark

<sup>‡</sup> Department of Chemistry and PULSE Institute, Stanford University, Stanford, CA 94305, USA, and SLAC National Accelerator Laboratory, Menlo Park, CA 94309, USA


## Introduction

### Motivation


- Polyenes like cyclopentadiene are ubiquitous as chromophores in diverse areas ranging from light harvesting and the vision process to devices such as light emitting diodes
- Polyenes possess low-lying optically dark states with large doubly excited character, which give rise to the observation of ultrafast excited state dynamics in the form of non-diabatic dynamics

### Focus

- Elucidate on structure-dynamics relationships focused on the influence of specific nuclear motion and substituents effects on dynamics at conical intersections
- Correlate dynamics simulations using Ab Initio Multiple Spawning<sup>[2]</sup> with experimental data from time-resolved mass spectrometry and photoelectron spectroscopy



The two  $\pi$ - and two  $\pi^*$ -molecular orbitals of cyclopentadiene from HOMO-1 to LUMO+1



Electronic spectrum of cyclopentadiene with indication of different transitions<sup>[1]</sup>

## Methods

### Ab Initio Multiple Spawning

- Nuclear time-dependent Schrödinger equation solved in a basis of gaussians (TBF)
- Electronic structure is calculated *on-the-fly*
- Parameters of basis functions  $\chi_j^l$  propagated according to Hamilton's equations
- Coefficients  $C_j^l(t)$  of wave function  $X_j(\mathbf{R}, t)$  propagated quantum mechanically
- New basis functions optimally spawned when coupling between surfaces is large

- Basis function :  $\chi_j^l(\mathbf{R}; \bar{\mathbf{R}}_j^l(t), \bar{\mathbf{P}}_j^l(t), \bar{\mathbf{V}}_j^l(t), \alpha_j^l) = \left(\frac{2\alpha_j^l}{\pi}\right)^{1/4} \exp\left[-\alpha_j^l(R_{ij} - \bar{R}_{ij}^l(t))^2 + i\bar{P}_{ij}^l(t)(R_{ij} - \bar{R}_{ij}^l(t))\right]$

$$\chi_j^l(\mathbf{R}; \bar{\mathbf{R}}_j^l(t), \bar{\mathbf{P}}_j^l(t), \bar{\mathbf{V}}_j^l(t), \alpha_j^l) = e^{i\bar{V}_j^l(t)} \prod_{\eta} \chi_{j\eta}^l(R_{\eta}; \bar{R}_{j\eta}^l(t), \bar{P}_{j\eta}^l(t), \alpha_{j\eta}^l)$$

- Wave function :  $X_j(\mathbf{R}, t) = \sum_j C_j^l(t) \chi_j^l(\mathbf{R}; \bar{\mathbf{R}}_j^l(t), \bar{\mathbf{P}}_j^l(t), \bar{\mathbf{V}}_j^l(t), \alpha_j^l)$


- Working eqs.  $\dot{\mathbf{S}}\mathbf{C} = -i(\mathbf{H} - i\dot{\mathbf{S}})\mathbf{C}$   $\mathbf{H}_{ij}^l = \langle \chi_i^l | \hat{\mathbf{H}} | \chi_j^l \rangle$   $\mathbf{S}_{ij}^l = \langle \chi_i^l | \chi_j^l \rangle \delta_{ij}$   $\dot{\mathbf{S}}_{ij}^l = \left\langle \chi_i^l \left| \frac{\partial}{\partial t} \right| \chi_j^l \right\rangle \delta_{ij}$

$$\frac{\partial \bar{R}_{j\eta}^l}{\partial t} = \frac{\bar{P}_{j\eta}^l}{m_{\eta}} \quad \frac{\partial \bar{P}_{j\eta}^l}{\partial t} = -\frac{\partial V_{j\eta}(\mathbf{R})}{\partial R_{j\eta}} \bigg|_{\bar{\mathbf{R}}_j^l(t)} \quad \frac{\partial \bar{V}_j^l}{\partial t} = -V_{j\eta}(\bar{\mathbf{R}}(t)) + \sum_{\eta} \frac{(\bar{P}_{j\eta}^l(t))^2}{2m_{\eta}}$$


### Initial Conditions

- 40 TBFs for CPD and 24 for Me<sub>4</sub>-CPD
- Positions and momenta sampled from the 0 K Wigner distribution of the harmonic approximation to the vibrational ground state


## Dynamics following Excitation to S<sub>1</sub>



Decay of S<sub>1</sub> population in CPD and Me<sub>4</sub>-CPD.




Scheme of the dynamics following excitation to S<sub>1</sub> as observed in the calculations.




### Bond Alternation

- Initial nuclear motion is primarily in-plane with no significant difference between the two molecules.




Time scale ~ 20 fs




### Backbone Distortion

- Out-of-plane distortion of the backbone is observed but the expectation value is similar in the two molecules for times > 50 fs.




~ 50 fs





### Double Bond Twist

- A significant slower torsion in the double bonds is observed for Me<sub>4</sub>-CPD compared to CPD.



~ 100 fs

## Electronic Structure



- MS-3-MR-CASPT2/6-31G\*\* with a (4,4) active space
- Analytic non-adiabatic couplings<sup>[3]</sup>
- Four Minimum Energy Conical Intersections (MECIs) located – three between S<sub>1</sub> and S<sub>0</sub> and one between S<sub>2</sub> and S<sub>1</sub>

- No minimum (zero gradient) located for neither S<sub>1</sub> nor S<sub>2</sub> as the minimum energy configuration located for both states is an MECI

### References


1. R. P. Frueholz et al., J. Chem. Phys., **1979**, 70, 2003-2013
2. M. Ben-Nun, T. J. Martinez, Adv. Chem. Phys., **2002**, 121, 439-512
3. T. Mori et al., J. Phys. Chem. A, **2012**, 116, 2808-2818

## Time-Resolved Photoelectron Spectra


- The dipole matrix element connecting the initial neutral state  $I$  and final cationic state  $F$  is evaluated in the single-active electron approximation using the strong orthogonality condition

$$\langle \Psi_I^N(\mathbf{r}_1, \dots, \mathbf{r}_N) | \hat{\mu}(\mathbf{r}_1, \dots, \mathbf{r}_N) | \Psi_F^{N-1}(\mathbf{r}_2, \dots, \mathbf{r}_N) \Psi_{kf}^e(\mathbf{r}_1) \rangle = \langle \phi_{IF}^D(\mathbf{r}_1) | \hat{\mu}(\mathbf{r}_1) | \Psi_{kf}^e(\mathbf{r}_1) \rangle_{\mathbf{r}_1}$$

$$\sigma_{\text{stick}}(E_k, \Delta t) \propto \sum_{I,F} \sum_f \left| \mathcal{E}_\omega(\Delta t) \cdot \langle \phi_{IF}^D(\mathbf{r}_1) | \hat{\mu}(\mathbf{r}_1) | \Psi_{kf}^e(\mathbf{r}_1) \rangle_{\mathbf{r}_1} \right|^2 \delta([\hbar\omega - \Delta E] - E_k)$$



Combined 1- and 2-photon spectrum for CPD. The band above 1 eV results from 2-photon ionization. Both observed bands are due to ionization out of the S<sub>1</sub> state.



1-photon spectrum for Me<sub>4</sub>-CPD. Both observed bands are due to ionization out of the S<sub>1</sub> state.